

May 2020

# Dark Energy and the Time Dependence of Fundamental Particle Constants

**Bodo Lampe**

II. Institut für theoretische Physik der Universität Hamburg  
Luruper Chaussee 149, 22761 Hamburg, Germany

## **Abstract**

The cosmic time dependencies of  $G$ ,  $\alpha$ ,  $\hbar$  and of Standard Model parameters like the Higgs vev and elementary particle masses are studied in the framework of a new dark energy interpretation. Due to the associated time variation of rulers, many effects turn out to be invisible. However, a rather large time dependence is claimed to arise in association with dark energy measurements, and smaller ones in connection with the Standard Model.

## I. Introduction

Dirac was one of the first to suggest that fundamental physical constants may vary in time due to the expansion of the universe[1]. Dirac concentrated on Newton's constant  $G$ , but since then a time dependence of  $c$ ,  $\alpha$ ,  $\hbar$  and so on has been considered possible as well ([2]-[18]).

From the 21st century perspective it is clear that if fundamental constants are time dependent in this way, the observed dark energy effect must have to do with it, because dark energy dominates the present expansion of the universe.

In the course of the present work **time dependencies** will therefore be partly reduced to a ('Planck' or 'dark' or cosmological) **energy dependence** of physical quantities and constants. This energy dependence is completely separate from and not to be confused with the usual energy dependence from the renormalization group.

The framework of the article will be the ordinary FLRW cosmology with a scale factor  $a(t)$  and a spatial curvature  $k$ , the latter assumed to be tiny (in accordance with observations). Furthermore, the so-called 'cosmic coordinate system' will be used, i.e. cosmic time  $t$  and proper distances  $r$  as parameters. This will prove to be optimal for the presentation.

It is well known that the fundamental spacetime constants  $c$ ,  $h$  and  $G$  can be used to define the Planck length, time and mass  $L$ ,  $T$  and  $M$  which describe the basic properties of space[m], time[s] and matter[kg]

$$L(t) = \sqrt{\frac{\hbar(t)G(t)}{c^3}} \quad T(t) = \sqrt{\frac{\hbar(t)G(t)}{c^5}} \quad M(t) = \sqrt{\frac{\hbar(t)c}{G(t)}} \quad (1)$$

One may invert these relations to obtain

$$c = \frac{L(t)}{T(t)} \quad \hbar(t) = E(t)T(t) \quad \kappa(t) = \frac{L(t)}{E(t)} \quad (2)$$

where  $E = Mc^2$  is the Planck energy and  $\kappa = G/c^4$  the Einstein constant.

A time dependence of these quantities has been anticipated here.  $t = 0$  is taken to be the present, so we have the present day values  $L_0 = L(0) =$  Planck length,  $T_0 = T(0) =$  Planck time and  $E_0 = E(0) =$  Planck energy. Numerical values are

$$L_0 = 1.6 \times 10^{-35}m \quad M_0 = 2.2 \times 10^{-8}kg \quad T_0 = 5.4 \times 10^{-44}s \quad (3)$$

No time dependence of  $c$  is indicated, because in the present model there is none - at least if one uses the above mentioned cosmic coordinates  $t$  and  $r$ , in which case the FLRW solution of the Einstein equations has the line element

$$ds^2 = -c^2 dt^2 + a(t)^2 \left[ \frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right] \quad (4)$$

with a constant i.e. time-independent speed of light.

$c$  being constant, one only needs to consider time dependencies of  $G$  and  $h$ .

Equivalently, since one has  $T(t)=L(t)/c$  one only needs to consider time dependencies of the Planck length  $L(t)$  and Planck energy  $E(t)$ .

Rewriting eq. (2) as

$$\hbar(t)c = E(t)L(t) \quad (5)$$

$$G(t) = c^4 L(t)/E(t) \quad (6)$$

one sees that there are 2 really fundamental time dependencies to be considered:

**-L(t)=the time dependence of the fundamental measure of space**

**-E(t)=the time dependence of the 'physically active' quantities - the 'quantities of motion', as Isaac Newton called them.**

Remark: The cosmic time dependence of elementary particle couplings like the fine structure constant  $\alpha$ , the elementary particle masses and so on is a different story. It will be treated in section V and will boil down to determine the time dependence of one other quantity:

**-J(t)=the time dependence of the 'internal exchange energy'** to be defined in section V.

## II. Measure-of-Space Equation

To determine  $L(t)$  and  $E(t)$  I introduce 2 equations:

$$\ddot{L} = -\frac{4\pi}{3}G\rho L - \omega^2(L - L_s) + \frac{\Lambda}{3}c^2 L \quad (7)$$

The idea behind this is that the universe is an elastic medium which consists of elementary constituents called tetrons[19, 20], and the bond length of these constituents is given by the Planck length  $L(t)$ , while the Planck energy  $E(t)$  measures the binding energy of every 2 bound constituents. It is to be noted that the




Figure 1: Schematic depiction of the universe as an elastic medium made of tiny constituents. Shown are the binding lines between nearest neighbour constituents. The 2 lattices represent the expanding universe at 2 times  $t_1 < t_2$ .

tetrons are invisible to us. All (ordinary and dark) matter particles and radiation we know are quasi-particles/wave-excitations of them and can propagate on the elastic medium.<sup>1</sup> We ourselves are wave-excitations, too, and because of this, the world appears Poincaré invariant to us, without a preferred rest system.

Since the medium is assumed to be elastic, in (7) a harmonic ansatz  $\ddot{L} = \dots - \omega^2(L - L_s)$  seems reasonable, cf. the discussion after eq. (11), in connection with (15) and at the end of this work. Within such a picture, in an expanding universe,  $L$  and  $E$  will vary with time (and so will  $h$  and  $G$  as well as all particle physics constants), and the next step is to make the most straightforward ansätze for these variations:

First of all, when the universe (=the elastic medium) expands, the variation of the Planck or bond length  $L(t)$  must reflect the general expansion as described by the FLRW expansion parameter  $a(t)$ . Eq. (7) relies on the simple assumption that on the average the bond length between 2 tetrons is always proportional to the scale parameter, i.e.  $a \sim L$  or equivalently

$$\frac{a(t)}{a_0} = \frac{L(t)}{L_0} \quad (8)$$

---

<sup>1</sup>In the tetron-model[19] our universe is embedded in a higher-dimensional space, and as an elastic medium it can thus acquire the full 3+1 GR curvature within this space, including the timely curvature related to expansion.

At first sight this equation may not seem reasonable to all readers. After all, the FLRW scale  $a(t)$  describes changes at cosmological distances, while the bond length  $L(t)$  between tetrons is microscopic in origin. The idea behind (8) is depicted schematically in Figure 1. The 2 lattices represent the expanding universe at 2 times  $t_2 > t_1$ .  $a(t)$  corresponds to the full extension of the lattice, while  $L(t)$  is the lattice spacing, i.e. the distance between 2 nearest neighbors. In Figure 1 the ratio  $a(t_2) : a(t_1)$  is given by 2. The same is true for the ratio  $L(t_2) : L(t_1)$ , because the average bond length between the lattice points grows in the same way as the lattice as a whole. This simple consideration is at the heart of (8) and also (through the FLRW equation) of (7).

Thus, the first term in (7) arises from the general relativistic deceleration of the universe through its matter content  $\rho$ , while the second term accounts for the dark energy phenomenon, however, not quite in the usual form of a cosmological constant (crossed out, because not utilized in the present work), but of a harmonic force  $-\omega^2(L - L_s)$ , that expands the elastic medium towards an equilibrium value  $L_s$  of the bond length  $L$ .

Eq. (7) tells us that linear forces are acting, one induced by (ordinary and dark) matter and driving the system towards  $L = 0$ , the other induced by the ('dark energy') tetron binding and driving it towards the equilibrium binding distance  $L_s$ . Presently we are in the region  $L_0 < L_s$ , so that  $-\omega^2(L - L_s)$  really is an expanding force. The value of  $\omega$  can and will be determined from a fit to dark energy measurements.

In the course of time, i.e. with increasing  $L$ , the matter force becomes smaller because the matter density dilutes according to  $\rho = \rho_0 L_0^3 / L^3$ . This is a well known effect and makes the first term on the RHS of (7) behave like  $\sim 1/L$  instead of  $\sim L$ .

The differential equation (7) can be solved using initial values

$$L(0) = L_0 \quad \dot{L}(0) = H_0 L_0 \quad (9)$$

where  $L_0$  is the (present day) value of the Planck length and  $H_0$  the Hubble constant (=present day value of the Hubble parameter  $H(t) = \dot{a}/a = \dot{L}/L$ ). The solution will be given later in (20).

From the initial conditions it is immediately clear that  $\omega$  is naturally of the order of

$H_0$ , In section IV this will be confirmed by fitting with observations.  $\omega$  and  $H(t)$  are extremely small frequencies corresponding to an approximately harmonic movement of the universe as a whole and a priori have little to do with the Planck frequency  $1/T_P$  which is the local response frequency of a single tetron in the elastic medium.

$$H_0 T_P \approx 1.18 \times 10^{-61} \quad (10)$$

So seemingly, there are 2 very different fundamental scales in the universe: one is the single tetron binding energy/Planck energy  $E$  and the other is the collective dark energy of the universe as a whole, which drives it to its equilibrium value. However, **this collective drive is just a reflection of the microscopic tetron binding energy having a minimum at bond length  $L_s$** . Therefore, although the values of  $E$  and  $H$  are vastly different, the time dependencies  $E(t)$  and  $H(t)$  are connected. See eq. (18) later.

In other words, due to the homogeneity of the elastic medium, the time behaviour of the microscopic tetron energy  $E(t)$  and that of the cosmological frequency  $H(t)$  can be related, as will be seen in eq. (18).

### III. Quantities-of-Motion Equation

If one thinks it over, a time dependent  $L(t)$  has long been observed, namely in the form of the cosmological redshift. Usually this time dependence is not put into  $L$ ,  $G$  or  $h$ , as in eqs. (5) and (6), but into the redshifted photon frequency  $f$  and the expansion parameter  $a$ . This is possible, because these quantities always appear in products  $h*f$  and  $G*a$ , respectively. So one can choose whether to absorb the time dependence of  $L$  in  $h$  and  $G$  or in  $f$  and  $a$ . The conventional choice is to keep  $G$  and  $h$  constant. We shall follow this choice - as far as the variation of the Measure-of-Space equation is concerned.

From this point of view, the ansatz of a time dependent  $L(t)$  is not so much new [apart from the modified cosmological constant approach to dark energy with  $-\omega^2(L - L_s)$  instead of a  $\Lambda$ -term].

As for the time dependence of the Planck energy  $E$ , the situation is different, i.e. there will be something new:




Figure 2: The binding energy  $E$  of 2 constituents as a function of their bond length  $L$ . At present  $t=0$  one has the Planck energy  $E_0$  and the Planck length  $L_0$ . The expansion of the universe through dark energy corresponds to the elastic bonds expanding towards equilibrium values  $E_s$  and  $L_s$ . In the neighbourhood of  $L_s$  the quadratic dependence  $E(L)$  of eq. (12) is a good approximation.

$E$  can be interpreted as the binding energy among the constituents of the elastic medium which is our universe. Not too far away from the equilibrium at  $L = L_s$  it can be expanded in a power series of  $(L - L_s)^2$

$$E(L) = C + D(L - L_s)^2 + O(L - L_s)^4 \quad (11)$$

where the terms of order  $(L - L_s)^4$  and higher will be neglected in the following. This corresponds to approximating the curve in figure 2 by a parabola in the neighborhood of  $L_s$ , which should be a reasonable approximation not too far away from the minimum at  $L = L_s$ .

The constants  $C$  and  $D$  can be determined from the conditions that  $E(L_0) = E_0$  and  $E(L_s) = E_s$ . One obtains

$$E(L) = E_s - (E_s - E_0) \left( \frac{L - L_s}{L_0 - L_s} \right)^2 = E_s \left[ 1 - \left( 1 - \frac{E_0}{E_s} \right) \left( \frac{1 - L/L_s}{1 - L_0/L_s} \right)^2 \right] \quad (12)$$

As will turn out, the energy difference  $E_0 - E_s$  triggers the harmonic dark energy term  $\sim \omega^2$  in eq.(7), i.e. the accelerated expansion of the universe.

The approximate behavior of  $E(L)$  is that of a parabola and together with the

solution  $L(t)$  to (7) one deduces the time dependence  $E(t)$  as needed in eqs. (5) and (6).

Since we have absorbed the time dependence of  $L$  into the redshift definition, one only has to consider time dependencies of  $\hbar$  and  $G$  through  $E(t)$ , cf. eqs. (5) and (6). These dependencies may be rewritten as

$$\hbar(t) = \hbar_0 \frac{E(L)}{E(L_0)} \quad (13)$$

$$G(t) = G_0 \frac{E(L_0)}{E(L)} \quad (14)$$

with  $E(L)[= E(L(t))]$  to be taken from (12).

Considered as a binding energy,  $E(t)$  is negative, i.e. (5) and (6) are better to be replaced by  $\hbar(t)c = |E(t)|L(t)$  and  $G(t)|E(t)| = c^4L(t)$  with  $|E| = Mc^2$  being the Planck energy. Since  $E(t)$  is negative and presently becomes more negative as it approaches its minimum value  $E_s$ , one concludes from eq. (13) that **Plancks constant presently goes up with time, whereas the gravitational coupling is decreasing.**

At this point one may worry, whether a varying  $E$  has a problem with energy conservation. Actually, this question also arises in connection with the redshift, and is usually answered by saying that energy 'goes into the metric'. Interpreting the universe as an elastic medium one can reformulate this by stating that energy goes into the total binding energy of the universe.

As discussed in connection with eq. (35), one can attribute the  $\omega$ -term in (7) to an 'energy'

$$W(L) = \frac{\omega_s^2}{2} - \frac{\omega^2}{2} \left( \frac{L - L_s}{L_0} \right)^2 = \frac{\omega_s^2}{2} \left[ 1 - \left( 1 - \frac{\omega_0^2}{\omega_s^2} \right) \left( \frac{1 - L/L_s}{1 - L_0/L_s} \right)^2 \right] \quad (15)$$

with

$$\omega^2 = \frac{\omega_s^2 - \omega_0^2}{(1 - L_s/L_0)^2} \quad (16)$$

Note the similarity between (12) and (15). Since the dark energy phenomenon is a smooth collective effect of all tetron binding energies  $E$  having a minimum at bond length  $L_s$ , i.e. the behaviour of  $W$  is a reflection of the tetron bond length driving towards its equilibrium value  $L_s$  (=the point where the tetron binding energy  $E$  is

having a minimum value  $E_s$ ), the time evolution of  $W$  and  $E$  is absolutely parallel, in an analogous way as the time evolution of  $a(t)$  is parallel to that of  $L(t)$ . In other words,  $E(L) \sim W(L)$  holds similarly as  $L(t) \sim a(t)$  for the cosmic scale factor  $a$  and the bond/Planck length  $L$ , cf. (8) and Figure 1, and one comes up with

$$\frac{W(L)}{W(L_0)} = \frac{E(L)}{E(L_0)} \quad (17)$$

The physical difference between  $W$  and  $E$  is that

- $E$  is the microscopic tetron binding energy and is roughly of the order of the present day Planck energy to be measured in Joule.
- the  $\omega$ 's are frequencies of the universe as a whole and measured in Hertz, and they are of the order of the Hubble parameter.

A direct consequence of (17) is

$$\frac{\omega_0^2}{\omega_s^2} = \frac{E_0}{E_s} \quad (18)$$

#### IV. Comparison with Astrophysical Data

In the laboratory it is more or less impossible to observe time variations of  $G$  and  $h$ , because via (1) these quantities define our rulers for mass and energy. While the universe expands, the rulers will expand, too.

In case of the redshift, astronomers were able to obtain relevant information on  $L(t)$  from observations of distant galaxies. In contrast, it seems difficult to measure the time variation (12) of energy from such observations, because any process, which took place in the past in some distant galaxy, will do so with the energy/rulers relations valid at that time, and when the produced particles arrive on earth they will interact with the detectors with the energy/rulers relations valid now; so that the observer will see no difference between processes now and then.

**As a consequence, time variations of  $h$  and  $G$  will generally not be visible.**

Not testable in particle processes, it turns out, however, that  $E(t)$  from eq. (12) can be directly observed in dark energy measurements. Dark energy observations do not usually concern the very early universe, so that the parabolic approximation

(12) should be good enough<sup>2</sup>. They are in effect testing eq. (7), and E(t) in (12) not only governs the  $\omega$ -term but according to (13) and (??) also enters the G-term on the RHS of (7).

In order to check this idea with astrophysical data, we go over from L(t) to the redshift z defined by

$$z(t) = \frac{a}{a_0} - 1 = \frac{L}{L_0} - 1 \quad (19)$$

The most precise measurement of the dark energy effect comes from the study of type-Ia supernovae in distant galaxies. I shall compare my redshift prediction to those data in a small-t approximation. This is justified because on cosmic scales the times involved are not too large.

Under this condition, up to  $O(t^4)$ , the solution to (7) can be written as

$$\begin{aligned} z = tH_0 &+ \frac{t^2 H_0^2}{2} \left[ -\frac{\Omega_M^0}{2} + \underbrace{\frac{\omega_0^2 \frac{E_s}{E_P} - 1}{H_0^2 \frac{L_s}{L_0} - 1}}_{\text{underwaves}} + \frac{\cancel{\Lambda c^2}}{3H_0^2} \right] \\ &+ \frac{t^3 H_0^3}{6} \left[ \underbrace{\Omega_M^0 \left( 1 + \frac{\frac{E_s}{E_P} - 1}{\frac{L_s}{L_0} - 1} \right)}_{\text{underlines}} - \underbrace{\frac{\omega_0^2 \frac{E_s}{E_P} - 1}{H_0^2 \left( \frac{L_s}{L_0} - 1 \right)^2}}_{\text{underwaves}} + \frac{\cancel{\Lambda c^2}}{3H_0^2} \right] \end{aligned} \quad (20)$$

The term indicated by underlines is the contribution from the time dependent Newton constant, the terms with underwaves come from the harmonic dark energy  $\omega$  contribution, and the crossed out terms from a cosmological constant (the latter to be ignored in the present model).

$$\Omega_M^0 = \frac{4\pi G_0 \rho_0}{3 H_0^2} \quad (21)$$

is the present day density parameter of matter in the universe, frequently used in this type of analysis. In the dark energy interpretation with a cosmological constant it comes out as roughly 0.3, which is usually considered a reasonable value.

As for any parabola, hidden in the parabolic dark energy (12) and (15) are 3 parameters, which need to be determined from observations. They may be chosen as

---

<sup>2</sup>For considering time variations of h and G in the very early universe, an approximation of the form (12) is not sufficient, because at small bond length L a typical binding energy is expected to be governed by a power behaviour of the form  $E(L) \sim L^{-n}$ .

- (i)  $\frac{E_s}{E_0} = \frac{\omega_s^2}{\omega_0^2} > 1$  = the ratio of the Planck energies resp dark energies at cosmic equilibrium and at present
- (ii)  $\frac{L_s}{L_0} > 1$  = the ratio of the tetron binding lengths at cosmic equilibrium and at present
- (iii)  $\frac{\omega_0^2}{H_0^2}$  = the ratio of the present dark energy over the present value of the Hubble constant.

Since there are more parameters than in the ansatz of a cosmological constant, the observations will only give relations between i, ii and iii. Furthermore, an estimate for  $\Omega_M^0$  has to be taken from other sources. Nevertheless, our next aim is to see what the observations allow to say.

A fit to the redshifts of supernovae yields[24]

$$z = tH_0 + \frac{t^2 H_0^2}{2}(1.00 \pm 0.05) + \frac{t^3 H_0^3}{6}(0.54 \pm 0.05) \quad (22)$$

Comparing with (21) one finds that it is easy to accommodate the data with the help of the quantities i, ii and iii. For example, choosing  $\Omega_M^0 = 0.3$  and

$$-L_s = 10L_0 \text{ one obtains } E_s = 1.34E_0 \text{ and } \omega_0^2 = 3.4H_0^2$$

$$-L_s = 2L_0 \text{ one obtains } E_s = 5.6E_0 \text{ and } \omega_0^2 = 0.25H_0^2$$

At first sight, the fact that data can be fitted this way so easily, seems to be a big surprise. After all, we are fitting numbers which usually are explained with an exponential increase due to a cosmological constant. The essential feature here is the contribution from the time variation of Newton's constant (underlined) which in combination with the harmonic dark energy contribution (underwaved) leads to an agreement with observations. The point is that since  $G(t)$  is going down with time, the retarding effect of (ordinary and dark) matter becomes smaller, and no exponential increase of the dark energy term as in the cosmological constant approach is needed.

In other words, although the harmonic force ansatz corresponds to a more moderate re-acceleration of the universe than the cosmological constant term, this is compensated by the time variation of energy as a whole which affects Newton's constant.

## V. Cosmic Time Dependence of Particle Physics Parameters

The analysis will now be extended to the 'constants', which describe the particle

physics interactions. All parameters of the Standard Model (SM) of particle physics will be considered, i.e.

-the 3 dimensionless gauge couplings: the weak and electromagnetic fine structure constants  $\alpha_{weak}$  and  $\alpha$  together with the QCD scale parameter  $\Lambda_{QCD}$ .

-2 further parameters of the electroweak Standard Model, in order to describe the Higgs potential. These will be chosen to be the Higgs mass  $m_H$  and the vacuum expectation value  $v$  of the Higgs field.

-the Yukawa couplings, which are all proportional to  $v$ .

Except for  $\alpha_{weak}$  and  $\alpha$ , all these parameters have dimension of energy. If one looks at the definition of the fine structure constant

$$\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} \quad (23)$$

it is the only dimensionless combination which can be built from the quantities  $e^2/\epsilon_0$ ,  $\hbar$  and  $c$ . As dimensionless, it is independent of the choice of rulers for time, length and energy<sup>3</sup>. This is good news, because in looking for a cosmic time dependence of  $\alpha$  one circumvents all the problems usually encountered in determining the time dependence of dimensionful quantities like  $E(t)$ . The *bad news* in considering ratios like  $\alpha$  is that most effects tend to drop out between numerator and denominator (see later).

An interesting point is that although  $\alpha$  itself is not an energy, it can be written as a ratio of forces or energies. Namely one can rewrite (23) as

$$\alpha = \frac{e^2}{4\pi\epsilon_0 r^{(2)}} / \frac{G_0 M_0^2}{r^{(2)}} \quad (24)$$

i.e. as the ratio of the electrostatic Coulomb (force) energy and the gravitational (force) energy of 2 point particles with elementary charge  $e$  and Planck mass  $M_0$  at an arbitrary distance  $r$ .

From this point of view the gravitational force is by no means small as compared to the electric force, but - for such tetron-like test particles - is 137 times stronger!

---

<sup>3</sup>The dependence on the Planck/tetron binding energy  $E$  is not to be confused with the Wilsonian running of coupling constants, i.e. the dependence of  $\alpha$  on the energies of particles in a scattering process.

The key relation here is

$$\hbar c = G_0 M_0^2 = L_0 E_0 \quad (25)$$

which follows from (2).

Defining  $Q^2 = e^2/[4\pi\epsilon_0]$  and introducing time dependencies, one has

$$\alpha(t) = \frac{Q^2(t)/L(t)}{E(t)} \quad (26)$$

where  $Q^2$  comprises the electromagnetic effect in a measurement-system independent way. Obviously,  $Q^2$  has the dimension of length $\times$ energy. Since measurements and astrophysical observations show almost no time variation of  $\alpha$ , the time dependence of  $Q^2/L$  must be the same as that of  $E(t)$  to a very good approximation.

Referring once again to the tetron model, this has to do with the fact, that the time dependence of  $Q^2$  is determined by that of the binding energy  $E(t)$ [19], so that any time dependence of  $\alpha$  drops out between numerator and denominator in (26).

To understand this point in detail, one should note that the tetron model is more than a microscopic theory for the cosmic elastic medium. The tetrans actually appear in the form of tetrahedrons which extend into a 3-dimensional internal space and whose excitations can be shown to represent the complete 3-family quark and lepton spectrum[19, 20].

The internal interactions among tetrans are typical quantum interactions in the sense that one always has 'exchange' energies in addition to 'direct' energies, simply because for 2 (or more) identical particles - tetrans in this case - with single wave functions  $f_1$  and  $f_2$  their total wave functions are either symmetric or antisymmetric of the form  $f_1(x_1)f_2(x_2) \pm f_1(x_2)f_2(x_1)$ . Correspondingly, the relevant 2-point function of the tetron Hamiltonian can be described as the sum of the Planck(=binding) energy  $E(t)$  and a function  $J(t)$  usually called the exchange energy. In the present case it may be called 'internal exchange energy' because it arises as an integral including the internal space, in which the tetrahedrons are living.

$$E = \int d^6x_1 \int d^6x_2 f_1(x_1)f_2(x_2)V(1-2)f_1(x_1)f_2(x_2) \quad (27)$$

$$J = \int d^6x_1 \int d^6x_2 f_1(x_1)f_2(x_2)V(1-2)f_1(x_2)f_2(x_1) \quad (28)$$

where the integrals are actually 6-dimensional, because they extend over both internal and physical space.  $V(1-2)$  is the potential between 2 tetrons with wave functions  $f_1$  and  $f_2$ .<sup>4</sup>

In a 6-dimensional environment the Green's function of the Laplace operator is  $r^{-4}$ , instead of  $r^{-1}$  in the 3-dimensional case. Therefore, the most promising choice seems to be

$$V(1-2) = \frac{N}{|x_1 - x_2|^4} \quad (29)$$

with some coupling constant  $N$ . A rough estimate of  $N$  can be obtained by equating  $V(1-2)$  at the Planck length to the Planck energy. This gives a value for the fundamental tetron coupling  $N$ :

$$\frac{N}{L_0^4} \approx E_0 \quad \implies \quad N \approx 10^{-130} \frac{m^6 kg}{s^2} \quad (30)$$

When trying to calculate  $E$  and  $J$  according to (27) and (28), one naturally runs into the so-called hierarchy problem of physics. Namely the question, why the relevant energy scales of gravity ( $E_0 \approx 10^{19} \text{GeV}$ ) and of particle physics ( $J_0 = 1 - 100 \text{GeV}$ ) are so much different. In the framework of the tetron model, the question can be reformulated: why is the exchange energy  $J$  so much smaller than the direct energy  $E$ ?

Looking at (27) and (28), one sees that  $J \ll E$  can happen, if the tetron wave functions are strongly localized. In the extreme case of delta functions one even finds, that the exchange integral vanishes, while the direct integral attains the value (30). Such an extreme localization is of course unnatural. In order to get  $J \approx 10^{-17} E$ , it is enough to demand that  $f(x)$  drops from its maximum value at  $x = 0$  by about

---

<sup>4</sup>If one looks into the details of the tetron model[19], the situation is a bit more complicated than described here. First of all,  $f_1$  and  $f_2$  are the wave functions of tetron-antitetron *pairs*, and  $V(1-2)$  is the potential between these 2 pairs. Secondly, to really calculate  $E$  and  $J$  from the 6-dimensional integrals one has to take the configuration of 2 adjacent tetrahedrons with at least 8 tetrons into account. Furthermore, there are actually 2 types of exchange integrals, one corresponding to the *inter-tetrahedral* interactions, which gives rise to the Fermi scale and is responsible for the large masses  $m_t$ ,  $m_W$  and  $m_H$  of order 100 GeV, and another one corresponding to the *inner-tetrahedral* interactions, which gives rise to the lighter fermion masses and the QCD scale.

a factor of 10 at  $x = L_0$ . This is because  $J$  is a multidimensional integral and to integrate the product  $dx_2 f_2(x_2) V(1-2) f_1(x_2)$  will give a suppression factor of roughly  $\sim 0.1$  for each of the 6 dimensions. Similarly for the  $x_1$ -integration.

Except for  $\alpha$ , which is constant, I will argue that  **$J(t)$  gives a universal time dependence for all internal/particle interactions in a similar way as does the Planck energy  $E(t)$  for the spacetime quantities of motion.** In other words, while the time-dependence of all dimensionful spacetime quantities is dictated by  $E(t)$ , the time dependencies of dimensionful SM particle properties like  $v$ ,  $m_H$ ,  $m_W$  and all quark and lepton masses can be described in terms  $J(t)$ .

To see how this works in detail, one should relate  $J$  to the electroweak symmetry breaking scale. This was already done in [19], where it was shown that the critical energy of the electroweak phase transition is given by an exchange integral  $J$  of the form (28). This is because in the tetron model the electroweak phase transition corresponds to an alignment of the tetrahedrons in the internal spaces, and the Curie energy of this phase transition is given by  $J$ . Since the critical energy of the electroweak phase transition is approximately given by the Higgs vev  $v$ , one has  $v = J$ .

It is well known that all particle masses in the SM are proportional to  $v$ . Therefore,  $J$  enters all dimensionful parameters of the electroweak SM - the fermion masses, the Higgs vev and the masses of the weak gauge bosons - in a linear way. All these quantities are  $\sim J(t)$ .

Just as for  $E$ , the time-dependence of  $J$  arises through the time variation of the bond length  $L(t)$ , i.e. through the expansion of the universe. If one would calculate the integrals  $J$  and  $E$  (27) and (28) as a function of  $L$ , then knowing  $L(t)$  according to (7), one could deduce from that the time-dependence of  $v$ , of the Fermi constant (??) and of all other parameters, and compare it to present upper limits[25, 2].

Unfortunately, the situation is not that simple. First of all, as mentioned in footnote 4, the integrals are difficult to calculate. Secondly, in everything we do, in every experiment we undertake, we encounter the Planck energy  $E(t)$  as a ruler, whose time dependence influences our perception of dimensionful quantities like  $v$ ,  $G_F$ ,  $m_W$  and so on. To say it plainly, **the time dependence we can perceive is not that of  $J(t)$  but that of the ratio  $J(t)/E(t)$ .**

This means: if we consider, for example, a matter particle with mass  $m_0$  in the present epoch, our perception of the time development  $m(t)$  of  $m_0$  does not follow<sup>5</sup>

$$m(t) = m_0 \frac{J(t)}{J_0} \quad (31)$$

but

$$m(t) = m_0 \frac{J(t)/J_0}{E(t)/E_0} \quad (32)$$

In the ideal case, that J and E would have an identical time dependence, the time dependence of m or of a dimensionful SM parameter like (??) could never be measured.

By analyzing the structure of the direct and the exchange integrals E and J in some detail, one can indeed show, that their dependence on the bond length L is quite similar, both with an extremum at nearly the same value  $L_s$ . Making an ansatz for J(L) analogous to that for E(L) in (12)

$$J(L) = J_s - (J_s - J_0) \left( \frac{L - L_s}{L_0 - L_s} \right)^2 = J_s \left[ 1 - \left( 1 - \frac{J_0}{J_s} \right) \left( \frac{1 - L/L_s}{1 - L_0/L_s} \right)^2 \right] \quad (33)$$

one sees that the crucial part is the ratio  $J_0/J_s$ . To the extent that the equality

$$\frac{J_s}{J_0} = \frac{E_s}{E_0} \quad (34)$$

holds, a time dependence of SM parameters cannot be measured. Conversely, any observed time dependence in a SM parameter can be traced back to a deviation from (34).

The integrals J and E can be analysed on a qualitative level, and according to this analysis the relation (34) is approximately true. On the other hand there is no a priori reason, why it should be exactly true. First of all, the integrals (27) and (28) are definitely distinct. Secondly, particle physics interactions have to do with inner symmetries not contained in the energetic analysis of the elastic universe [governed by E(t)]. Therefore, although present observations only give upper limits on time dependencies of SM parameters, their cosmic time dependence at least in principle follows its own rule, given by J(t).

---

<sup>5</sup>Note there is no problem with the principle of equivalence because the heavy mass and the inert mass are both developing with J(t).

## VI. Discussion

In this study a theory concerning the time dependence of all known fundamental physical parameters has been developed. It rests on the idea that dark energy is a harmonic rather than an exponential effect, which is furthermore related to the binding energy of the underlying constituents of the universe. As has been shown, one is led to a time-dependence of Newton's and Planck's constant. These effects, however, are usually impossible to measure - except in the dark energy itself and in certain particle physics properties.

Furthermore

-microscopic ( $L$ ) und cosmic ( $a$ ) length scales are connected in a simple linear kind of way ('the universe expands in the same manner as the tetron bonds expand'), cf. eq. (8) and Figure 1.

-In an analogous fashion, Planck energies  $E(t)$  and dark energies  $\omega$  are linearly related ('the total dark energy of the universe increases proportional to the single tetron binding energy') via (17).

Since the universe is rather cool by now and apparently expands in a rather homogeneous way, these assumptions are expected to be very good approximations. This expectation was substantiated in section IV by proving that it leads to agreement with present day dark energy observations. Thereby it has turned out that there is a significant contribution to the observed dark energy effect from the time variation of Newton's constant. Since  $G(t)$  is going down with time, the retarding effect of ordinary matter becomes smaller, and no exponential increase of the dark energy effect as in the cosmological constant approach is needed.

The Planck energy  $E_0$  and its time-dependent generalization  $E(t)$  play a central role in the considerations presented here, see (5), (6) and (12). Actually,  $E_0$  has been used in this paper with 2 meanings:

-it represents the gravitational energy of the interaction of 2 matter particles with Planck mass  $M_0$  at Planck distance  $L_0$ , i.e.  $E_0 = G_0 M_0^2 / L_0$ .

-it describes the binding energy of 2 tetrons bound at distance  $L_0$ .

Concerning the fundamental parameters of particle physics, it was shown that they depend on cosmic time via the internal exchange function  $J$ , whose dependence on  $L$

is similar but not exactly the same as that of E. With the advent of higher precision observations, this effect may become observable.

A remaining problem is the calculation of E(t) (Planck energy) and J(t) (internal exchange energy) from first principles, i.e. from fundamental tetron interactions. In principle, this can be done using eqs. (27) and (28), as soon as the precise form of the fundamental tetron interactions is known.

Finally, I want to discuss the question of energy conservation in the present framework. First of all, it should be stressed that energy is conserved locally in the interplay of processes between tetrans and ordinary and dark matter, just as it is in GR in the interplay between the metric, the vacuum energy and ordinary and dark matter.

The possibility that the observed acceleration of the Universe and the possible time variation of the fundamental constants are two manifestations of the same underlying dynamics ('micro and macro connection'), has already been considered in a somewhat different context by the authors of [21, 22, 23]. In those papers a cosmic time dependent vacuum energy density has been introduced, and it was nicely shown how energy conservation can be maintained by an interplay between matter and the 'running vacuum'.

There is some similarity of this approach to the present work, however the arguments of [21, 22, 23] are not directly applicable here, because the conserved energy of the elastic tetron medium cannot be given in a Lorentz invariant way. Actually, the conserved quantity, from which the vibrational  $\omega^2$ -term in eq. (7) can be derived is given by  $\dot{L}^2 + \omega^2(L - L_s)^2$ , or, using (8), by

$$\frac{1}{2}\dot{a}^2 + \frac{\omega^2}{2}(a - a_s)^2 = const \quad (35)$$

for a matter free elastic universe.  $a_s$  is the equilibrium value of  $a(t)$ , at which the tetron binding energy has a minimum, cf. figure 2. The  $\omega^2$ -term is not a part of the Einstein theory. It represents the energy of the elastic medium universe which vibrates with a 'breathing frequency'  $\omega$  around  $a = a_s$ . Furthermore, this expression is valid only in the FLRW cosmic coordinate system and therefore coordinate dependent and the model cannot be presented in a Lorentz covariant form.

The fact that it cannot be made a Lorentz covariant part of the Einstein theory

should not be considered as a surprise, because general relativity is a theory of local curvature induced by energy-momentum and does not know about the equilibrium of the underlying elastic medium at  $a = a_s$ . In the elastic picture of the universe, the FLRW cosmic coordinate system is a preferred system, where the tetrons do not move except for their general elastic expansion as in figure 1. Ordinary and dark matter particles are quasi-particle waves, gliding on the tetron substrate. Since they respect Lorentz covariant wave equations, their physics looks the same in all inertial coordinate systems. The elastic tetron background, on the other hand, should be described on its preferred FLRW 'rest system' figure 1, and for this the energy principle (35) can be formulated.

## References

- [1] P. A. M. Dirac, Nature 139, 323 (1937); Proc. Roy. Soc. A165, 199 (1938).
- [2] see the review by J.P. Uzan, Living Reviews in Relativity 14 (2011) 2.
- [3] M. J. Duff, L. B. Okun and G. Veneziano, JHEP 0203, 023 (2002).
- [4] J. W. Moffat, Int. J. Mod. Phys. D2, 351 (1993).
- [5] J. D. Barrow, Phys. Rev. Lett. 82 (1999) 884.
- [6] J. D. Bekenstein, J. D. Phys.Rev. D25 (1982) 1527.
- [7] H. Fritzsche and J. Sola, Adv. High Energy Phys. 2014 (2014) 361587.
- [8] J. Magueijo, Rept. Prog. Phys. 66 (2003) 2025.
- [9] J. K. Webb et al. Phys.Rev.Lett. 87 (2001) 091301.
- [10] J. A. Belinchon et al., Int. J. Mod. Phys. D12 (2003) 1113.
- [11] J. C. Carvalho et al. Phys. Rev. D46 (1992) 2404.
- [12] T. Chiba, Prog. Theor. Phys. 126 (2011) 993.
- [13] X. Calmet et al., Phys. Lett. B540 (2002) 173.

- [14] C. L. Gardner, Phys. Rev. D68 (2003) 043513.
- [15] D. F. Mota et al. Phys.Lett. B581 (2004) 141.
- [16] D. Parkinson et al. Phys. Lett. B578 (2004) 235.
- [17] E. J. Copeland, Phys. Rev. D69 (2004) 023501.
- [18] D. J. Shaw et al., Phys. Rev. D71 (2005) 063525.
- [19] for a review see B. Lampe, arXiv:1505.03477 [hep-ph].
- [20] B. Lampe, Int. J. Mod. Phys. A30 (2015) 1550026.
- [21] H. Fritzsche and J. Sola, Class. Quant. Grav. 29 (2012) 215002.
- [22] H. Fritzsche and J. Sola, Mod. Phys. Lett. A, Vol. 30, No. 22 (2015) 1540034.
- [23] H. Fritzsche, J. Sola and Rafael C. Nunes, Eur.Phys.J. C77 (2017) 193.
- [24] A.G. Riess et al., Astronomical Journal 116 (1998) 1009; S. Perlmutter et al., Astrophysical Journal 517 (1998) 565.
- [25] A. Ferrero and B. Altschul, arXiv:1008.4769v2 [hep-ph]